


AD-A154 883 ON THE TURNING LOZANGE OF CONSTANT SIDE WHOSE VERTICES 1/1.  
ALTERNATE BETWEEN. (U) WISCONSIN UNIV-MADISON  
MATHEMATICS RESEARCH CENTER I J SCHOENBERG APR 85  
UNCLASSIFIED MRC-TSR-2813 DAAG29-88-C-0041 F/G 12/1 NL



END

FORMED

ONE



MICROCOPY RESOLUTION TEST CHART  
NATIONAL BUREAU OF STANDARDS-1963-A

AD-A154 803

MRC Technical Summary Report #2813

ON THE TURNING LOZANGE OF CONSTANT SIDE  
WHOSE VERTICES ALTERNATE BETWEEN  
TWO FIXED CIRCLES OF A RING

I. J. Schoenberg

Mathematics Research Center  
University of Wisconsin—Madison  
610 Walnut Street  
Madison, Wisconsin 53705

April 1985

DTIC FILE COPY (Received April 10, 1985)

Approved for public release  
Distribution unlimited

Sponsored by

U. S. Army Research Office  
P. O. Box 12211  
Research Triangle Park  
North Carolina 27709

DTIC  
ELECTE  
JUN 11 1985

S D  
G

85 06 10 165

UNIVERSITY OF WISCONSIN - MADISON  
MATHEMATICS RESEARCH CENTER

ON THE TURNING LOZANGE OF CONSTANT SIDE WHOSE VERTICES  
ALTERNATE BETWEEN TWO FIXED CIRCLES OF A RING

I. J. Schoenberg

Technical Summary Report #2813

April 1985

ABSTRACT

This note was written to call attention to the remarkable paper [1] by W. L. Black, H. C. Howland and B. Howland. Their result recalls the classical theorems of Poncelet and Steiner from the golden age of Geometry. [1] starts with two circles  $\Gamma$  and  $\tilde{\Gamma}$  in  $R^3$  satisfying the qualitative condition (A) and assumes that there is a closed equilateral polygon

$$\Pi_{2n} = P_1 P_2 \dots P_{2n} \quad (n > 2),$$

having the sides  $P_k P_{k+1} = d$  ( $k = 1, 2, \dots, n$ ;  $P_{2n+1} = P_1$ ) such that all vertices  $P_{2k-1}$  are on  $\Gamma$ , and  $P_{2k}$  are on  $\tilde{\Gamma}$  ( $k = 1, \dots, n$ ). It is shown that  $\Pi_{2n}$  can be turned around with constant  $d$ , so as to return to its initial position. This note, independent of [1], settles the case when the circles  $\Gamma$  and  $\tilde{\Gamma}$  are coplanar and  $n = 2$ , when  $\Pi_4$  becomes a lozange. It is shown that  $d^2 = a^2 + b^2 - c^2$ , where  $a$  and  $b$  are the radii of  $\Gamma$  and  $\tilde{\Gamma}$ , respectively, and  $c$  is the distance between the centers of  $\Gamma$  and  $\tilde{\Gamma}$ .

AMS (MOS) Subject: 51M15, 53A17

Key Words: <sup>cent</sup> Elementary Geometry, Linkages.

Work Unit Number 1 - Applied Analysis



Dist	Avail and/or Special	By	Distribution/	Availability Codes	Justification	Unannounced	DTIC TAB	NTIS GRA&I	Accession For
						<input checked="" type="checkbox"/>	<input type="checkbox"/>	<input type="checkbox"/>	

Sponsored by the United States Army under Contract No. DAAG29-80-C-0041.

# SIGNIFICANCE AND EXPLANATION



*on a theorem about zig-zags between two circles.*

This note is to call attention to the remarkable paper [1] by

W. L. Black, H. C. Howland and B. Howland, and may be read independently of

[1]. It explains the construction (Fig. 1) of a linkage composed of four

equal bars  $\overset{\text{sub } 1}{P_1} \overset{\text{sub } 2}{P_2} = \overset{\text{sub } 2}{P_2} \overset{\text{sub } 3}{P_3} = \overset{\text{sub } 3}{P_3} \overset{\text{sub } 4}{P_4} = \overset{\text{sub } 4}{P_4} \overset{\text{sub } 1}{P_1} = d$ , joined at the points  $\overset{\text{sub } k}{P_k}$ , with the following property: With  $\overset{\text{sub } 1}{P_1}$  and  $\overset{\text{sub } 3}{P_3}$  moving in the circular groove  $\overset{T}{A}$ , and  $\overset{\text{sub } 3}{P_3}, \overset{\text{sub } 4}{P_4}$  moving in the circular groove  $\overset{T'}{A}$ , it is shown that the lozange  $\overset{\text{sub } 1}{P_1} \overset{\text{sub } 2}{P_2} \overset{\text{sub } 3}{P_3} \overset{\text{sub } 4}{P_4}$  can be turned around. *keywords include: see page -A-*

The responsibility for the wording and views expressed in this descriptive summary lies with MRC, and not with the author of this report.

ON THE TURNING LOZANGE OF CONSTANT SIDE WHOSE VERTICES  
ALTERNATE BETWEEN TWO FIXED CIRCLES OF A RING

I. J. Schoenberg

I am writing this note to call attention to the remarkable paper [1] by W. L. Black, H. C. Howland and B. Howland. Their result which recalls the two classical theorems of V. Poncelet and J. Steiner (see e.g. [2, Chapter 14]), may be described as follows.

Let  $\Gamma$  and  $\tilde{\Gamma}$  be two circles in the 3-dimensional space  $R^3$  with the property

(A) There exists a number  $d$  such that each point of either circle is at the distance  $d$  from exactly two points on the other circle

Should the circles  $\Gamma, \tilde{\Gamma}$  be coplanar, then (A) will be satisfied if we make the assumption

(B) The smaller circle encloses the center of the larger circle.

Furthermore, the authors assume that there is a  $d$  with the following property: There is an equilateral closed polygon

$$\Pi_{2n} = P_1 P_2 \dots P_{2n} \quad (n \geq 2) ,$$

having all its sides =  $d$ , such that

$$(1) \quad P_{2k-1} \in \Gamma, P_{2k} \in \tilde{\Gamma} \quad (k = 1, 2, \dots, n) .$$

Then the polygon  $\Pi_{2n}$  can be turned around, with all its sides =  $d$ , so that the zig-zag property (1) holds, and returned to its initial position.

The authors do not describe a way of determining the (constant) size  $d$  of the side of  $\Pi_{2n}$ .

The present note should be regarded as an appendix to [1] that solves the case when  $n = 2$  and the circles are coplanar. It can be read independently. We prove

Theorem 1. Let  $O$  and  $\tilde{O}$  be the centers of the coplanar circles  $\Gamma$  and  $\tilde{\Gamma}$ , respectively, and let

$$(2) \quad a = \text{radius of } \Gamma, \quad b = \text{radius of } \tilde{\Gamma}, \quad c = O\tilde{O}.$$

We assume that the assumption (B) is satisfied. Then there is a 1-parameter family of lozanges  $\Pi_4$ , of constant sides =  $d$ , having the zig property (1) for  $n = 2$ , with  $d$  given by the equation

$$(3) \quad d^2 = a^2 + b^2 - c^2.$$

Proof.




Figure 1

I. Let us first show that there is a 1-parameter family of lozanges  $\Pi_4 = P_1P_2P_3P_4$  each of which enjoys the zig-zag property (1) for  $n = 2$ . Afterwards, we will show that the side  $d$  of  $\Pi_4$  satisfies the equation (3) so that  $d$  has the same value for all members of the family. Referring to Fig. 1 let  $\Pi_4 = P_1P_2P_3P_4$  be a lozange such that  $P_1$  and  $P_3$  are on  $\Gamma$ , and  $P_2, P_4$  on  $\tilde{\Gamma}$ . Evidently  $P_1P_2P_3$  is isosceles and so is  $P_1\tilde{O}P_3$ . It follows that  $\tilde{O}$  is on the diagonal  $P_2P_4$ . Likewise  $P_4P_1P_2$  and  $P_4\tilde{O}P_2$  are isosceles triangle, and therefore  $\tilde{O}$  is on the diagonal  $P_1P_3$ . Moreover  $P_1P_3$  and  $P_2P_4$  are evidently perpendicular to each other.

This makes the construction of  $\Pi_4 = P_1P_2P_3P_4$  evident: Having picked  $P_1$  arbitrarily on  $\Gamma$ , we draw the line joining  $P_1$  to  $\tilde{O}$  and intersecting  $\Gamma$  again in  $P_3$ . Next we drop the perpendicular from  $\tilde{O}$  onto  $P_1\tilde{O}P_3$ . With  $A = P_1P_3 \cap P_2P_4$  we evidently have  $P_1A = AP_3$  and  $P_2A = AP_4$ , so that  $\Pi_4 = P_1P_2P_3P_4$  is a lozange having the side  $d = P_1P_2$ .

We recall that  $P_1$  was an arbitrary point of  $\Gamma$ ; the constancy of  $d$  will follow as soon as we establish the equation (3), for it shows that  $d$  does not depend on the choice of  $P_1$  on  $\Gamma$ .

II. Let us prove the equation (3). To simplify our notations, and referring to Fig. 1, we write

$$e = A\tilde{O}, f = \tilde{O}P_1, g = A\tilde{O}, h = \tilde{O}P_2.$$

The Pythagorean theorem gives the following equations

$$d^2 = (P_1P_2)^2 = (e+f)^2 + (g+h)^2 = e^2 + f^2 + g^2 + h^2 + 2(e f + g h),$$

$$a^2 = (e+f)^2 + g^2 = e^2 + g^2 + f^2 + 2ef,$$

$$b^2 = (g+h)^2 + e^2 = g^2 + h^2 + e^2 + 2gh,$$

$$c^2 = e^2 + g^2.$$

These show that  $d^2 + c^2 = a^2 + b^2$  and the equation (3) is established. The constancy of  $d$  immediately implies the turning property of the lozange  $\Pi_4$ .



Remarks. 1. On Fig. 1 we see that if  $0 = \tilde{0}$ , hence  $c = 0$  then  $a = AP_1$  and  $b = AP_2$ , and therefore the equation (3) may be regarded as a generalization of the Pythagorean theorem

$$(P_1P_2)^2 = (AP_1)^2 + (AP_2)^2 .$$

2. Our result would make an attractive linkage that would be fun to handle: The four equal side of  $\Pi_4$  should be linked at the vertices, while the two pairs of opposite vertices  $P_1, P_3$  and  $P_2, P_4$  should fit into the circular grooves  $\Gamma$  and  $\tilde{\Gamma}$ , respectively. From the general result of [1] we may similarly make a linkage from the closed equilateral polygon  $\Pi_{2n}$ , with  $P_{2k-1}$  fitting into the circular groove  $\Gamma$ , and  $P_{2k}$  into  $\tilde{\Gamma}$ . Clearly  $n = 3$ , or  $n = 4$ , are the most likely choices. For  $n > 2$  the side  $d$  must be well approximated graphically, by trial and error, after the grooves  $\Gamma$  and  $\tilde{\Gamma}$  have been chosen.

### References

1. W. L. Black, H. C. Howland and B. Howland, A theorem about zig-zags  
between two circles, American Mathematical Monthly, 81 (1974),  
754-757.
2. I. J. Schoenberg, Mathematical Time Exposures, Mathematical Association of  
America, 1982.

IJS/jvs

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER #2813	2. GOVT ACCESSION NO. AD-A154803	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle)  On the Turning Lozange of Constant Side Whose Vertices Alternate Between Two Fixed Circles of a Ring		5. TYPE OF REPORT & PERIOD COVERED Summary Report - no specific reporting period
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s)  I. J. Schoenberg		8. CONTRACT OR GRANT NUMBER(s)  DAAG29-80-C-0041
9. PERFORMING ORGANIZATION NAME AND ADDRESS Mathematics Research Center, University of 610 Walnut Street Wisconsin Madison, Wisconsin 53705		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS Work Unit Number 1 - Applied Analysis
11. CONTROLLING OFFICE NAME AND ADDRESS U. S. Army Research Office P.O. Box 12211 Research Triangle Park, North Carolina 27709		12. REPORT DATE April 1985
		13. NUMBER OF PAGES 5
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report)  UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report)  Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number)  Elementary Geometry, Linkages		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number)  This note was written to call attention to the remarkable paper [1] by W. L. Black, H. C. Howland and B. Howland. Their result recalls the classical theorems of Poncelet and Steiner from the golden age of Geometry. [1] starts with two circles $\Gamma$ and $\bar{\Gamma}$ in $R^3$ satisfying the qualitative condition (A) and assumes that there is a <u>closed equilateral</u> polygon $\Pi_{2n} = P_1 P_2 \dots P_{2n} \quad (n \geq 2) \quad ,$		

ABSTRACT (continued)

having the sides  $P_k P_{k+1} = d$  ( $k = 1, 2, \dots, n$ ;  $P_{2n+1} = P_1$ ) such that all vertices  $P_{2k-1}$  are on  $\Gamma$ , and  $P_{2k}$  are on  $\tilde{\Gamma}$  ( $k = 1, \dots, n$ ). It is shown that  $\Pi_{2n}$  can be turned around with constant  $d$ , so as to return to its initial position. This note, independent of [1], settles the case when the circles  $\Gamma$  and  $\tilde{\Gamma}$  are coplanar and  $n = 2$ , when  $\Pi_4$  becomes a lozange. It is shown that  $d^2 = a^2 + b^2 - c^2$ , where  $a$  and  $b$  are the radii of  $\Gamma$  and  $\tilde{\Gamma}$ , respectively, and  $c$  is the distance between the centers of  $\Gamma$  and  $\tilde{\Gamma}$ .

**END**

**FILMED**

**7-85**

**DTIC**